

Protein Science

Crystallographic studies of the complexes of antiviral protein griffithsin with glucose and N-acetylglucosamine

Natasza E. Ziólkowska, Shilpa R. Shenoy, Barry R. O'keefe and Alexander Wlodawer

Protein Sci. published online Jun 13, 2007;
Access the most recent version at doi:[10.1110/ps.072889407](https://doi.org/10.1110/ps.072889407)

P<P	Published online June 13, 2007 in advance of the print journal.
Email alerting service	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here

Notes

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To subscribe to *Protein Science* go to:
<http://www.proteinscience.org/subscriptions/>

PROTEIN STRUCTURE REPORT

Crystallographic studies of the complexes of antiviral protein griffithsin with glucose and N-acetylglucosamine

NATASZA E. ZIÓŁKOWSKA,¹ SHILPA R. SHENOY,² BARRY R. O'KEEFE,³
AND ALEXANDER WLODAWER¹

¹Protein Structure Section, Macromolecular Crystallography Laboratory, National Cancer Institute, NCI-Frederick, Frederick, Maryland 21702-1201, USA

²Molecular Targets Development Program, SAIC-Frederick, Inc., National Cancer Institute, NCI-Frederick, Frederick, Maryland 21702-1201, USA

³Molecular Targets Development Program, Center for Cancer Research, National Cancer Institute, NCI-Frederick, Frederick, Maryland 21702-1201, USA

(RECEIVED March 21, 2007; FINAL REVISION March 30, 2007; ACCEPTED April 2, 2007)

Abstract

Crystal structures of complexes of an antiviral lectin griffithsin (GRFT) with glucose and N-acetylglucosamine were solved and refined at high resolution. In both complexes, all six monosaccharide-binding sites of GRFT were occupied and the mode of binding was similar to that of mannose. In our previous attempts to obtain a complex with N-acetylglucosamine by soaking, only a single site was occupied; thus, cocrystallization was clearly superior despite lower concentration of the ligand. Isothermal titration calorimetric experiments with N-acetylglucosamine, glucose, and mannose provided enthalpic evidence of distinct binding differences between the three monosaccharides. A comparison of the mode of binding of different monosaccharides is discussed in the context of the antiviral activity of GRFT, based on specific binding to high-mannose-containing complex carbohydrates found on viral envelopes.

Keywords: lectins; monosaccharide binding; antiviral activity; cocrystallization

Griffithsin (GRFT) is a potent antiviral lectin isolated from the red alga *Griffithsia* sp. (Mori et al. 2005). GRFT has been reported to exhibit significant activity against human immunodeficiency virus (HIV) and the coronavirus that causes severe acute respiratory syndrome (SARS) (Ziółkowska et al. 2006). The antiviral activity of GRFT is due to its binding to high-mannose oligosaccharides on viral envelope glycoproteins. Surprisingly for a protein with such potent anti-HIV activity ($EC_{50} = 0.043\text{--}1.0$ nM), the binding of GRFT to soluble HIV gp120 was inhibited by mannose, glucose, and N-acetylglucosamine (Mori et al.

2005). Generally, functionally similar monosaccharide-specific lectins (Ziółkowska and Wlodawer 2006) have been reported to have anti-HIV activity at concentrations ~1000-fold higher than that reported for GRFT (Charan et al. 2000). The unique potency to GRFT makes further study of its specific monosaccharide interactions of continuing interest.

Crystal structures of carbohydrate-free GRFT, as well as of its complexes with monosaccharides and disaccharides such as mannose, N-acetylglucosamine, $1 \rightarrow 6\alpha$ -mannobiose, and maltose were solved and refined at high resolution (Ziółkowska et al. 2006; 2007). The fold of GRFT corresponds to the β -prism-I motif (Chothia and Murzin 1993), which consists of three repeats of an antiparallel, four-stranded β -sheet that form a triangular prism. The first two β -strands of one chain are associated with 10 strands of the other chain and vice versa; thus, GRFT can be described

Reprint requests to: Alexander Wlodawer, Macromolecular Crystallography Laboratory, NCI, Frederick, MD 21702-1201 USA; e-mail: wlodawer@ncifcrf.gov; fax: (301) 846-6322.

Article published online ahead of print. Article and publication date are at <http://www.proteinscience.org/cgi/doi/10.1110/ps.072889407>.

as a domain-swapped dimer (Ziółkowska et al. 2006). A monomer of GRFT contains three almost identical carbohydrate-binding sites, each capable of binding a monosaccharide; thus, six principal sites are found in the dimer. Binding of oligosaccharides to multiple sites on a GRFT molecule provides the basis of its unique antiviral activity, which is >1000-fold more potent than the anti-HIV activities reported for several monosaccharide-specific lectins from the same structural family (Charan et al. 2000; Ziółkowska et al. 2006). This potency makes GRFT a promising candidate for development as a pharmaceutical agent for prevention of viral infection, specifically in the form of a female-controlled anti-HIV microbicide.

We present here the structures of GRFT cocrystallized with glucose and N-acetylglucosamine. No structure of glucose-bound GRFT had been reported, whereas the previously determined structure of the complex with N-acetylglucosamine (Protein Data Bank [PDB] code 2gue) was obtained by soaking, with only one binding site occupied by the carbohydrate.

Results and Discussion

Superposition of the complex of GRFT with glucose on the coordinates of the mannose complex of GRFT with mannose refined at 1.78 Å resolution (PDB code 2guc) (Ziółkowska et al. 2006) yielded a root mean square

(RMS) difference of 0.41 Å for the whole dimer. The largest differences were observed for Ser1A, Leu2A, Tyr121A, Ser54B, and Gly55B. An analogous superposition of the complex with N-acetylglucosamine yielded a RMS difference of 0.43 Å for the whole dimer. The largest differences were observed for Ser1A, Leu2A, Ser13B, Ser54B, and Gly55B.

GRFT was cocrystallized with glucose and N-acetylglucosamine in space group P2₁2₁2₁ (Table 1), isomorphous to the unliganded form (PDB code 2gty) that was previously used for soaking with N-acetylglucosamine (PDB code 2gue) but different from the cocrystals with mannose, maltose, and 1 → 6α-mannobiose. Nevertheless, the mode of sugar binding appears the same in all cocrystal forms of monosaccharides (Fig. 1). All six binding sites are occupied, and the conformation of the sugar molecules is similar to that of mannose. The pattern of hydrogen bonding in sugar binding sites for all cocrystal complexes of GRFT is also well conserved.

The difference between molecules of glucose and mannose is in the orientation of the O1 atom, with the details of the hydrogen bonding pattern of a glucose molecule shown in Figure 2A. The difference in sugar conformation between these monosaccharides does not appear to affect interactions with GRFT, since O1 points away from the protein chain. Similarly, the acetyl group of N-acetylglucosamine is directed away from the protein

Table 1. Statistics of data collection and structure refinement

	Crystal	
	Complex of GRFT with glucose	Complex of GRFT with N-acetylglucosamine
Data collection		
Space group	P2 ₁ 2 ₁ 2 ₁	P2 ₁ 2 ₁ 2 ₁
Cell parameters	a = 37.49 Å, b = 53.67 Å, c = 103.83 Å; α = β = γ = 90°	37.57 Å, 53.41 Å, 104.01 Å, α = β = γ = 90°
Molecules/a.u.	2	2
Resolution (Å)	30–1.50	30–1.56
Total reflections	882,464	477,222
Unique reflections	30,820	30,231
Completeness (%) ^a	89.6 (97.3)	98.8 (97.3)
Avg. I/σ	17.1 (7.2)	17.3 (1.95)
R _{merge} (%) ^b	8.0 (44.0)	8.4 (69.8)
Refinement statistics		
R (%) ^c	15.71	16.51
R _{free} (%) ^d	20.35	20.51
RMS deviations bond lengths (Å)	0.021	0.021
RMS deviations angles (°)	2.1	1.9
PDB code	2nuo	2nu5

The number of unique reflections is given first with Friedel pairs unmerged, and after merging in parentheses.

^aThe values in parentheses relate to the highest resolution shell.

^bR_{merge} = $\sum |I - \langle I \rangle| / \sum I$, where I is the observed intensity and $\langle I \rangle$ is the average intensity obtained from multiple observations of symmetry-related reflections after rejections.

^cR = $\sum ||F_o| - |F_c|| / \sum |F_o|$, where F_o and F_c are the observed and calculated structure factors, respectively.

^dR_{free} = defined by Brünger (1992).




Figure 1. Superposition of the structures of mannose (green) and glucose (yellow, *A*), or N-acetylglucosamine (white, *B*) in the binding sites 1–3 in the cocrystal complexes of GRFT. The coordinates of only the predominant α anomer of N-acetylglucosamine are shown.

(Fig. 2B). In isothermal titration calorimetry binding experiments with GRFT, interaction heats were detected for both glucose and N-acetylglucosamine. These heats of interaction were comparable to that of mannose (Fig. 3), with a 1.5-fold decrease for glucose and a twofold decrease for N-acetylglucosamine (Table 2). In binding experiments with other monosaccharides to GRFT, only minimal or no heats of interaction were detected (data not shown). However, a subtle but important difference between mannose and either glucose or N-acetylglucosamine was detected in the calorimetry binding experiments. As was previously reported (Ziółkowska et al. 2007), binding between mannose and GRFT was fully saturable ($K_d = 102 \mu\text{M}$). In the case of glucose and N-acetylglucosamine, although interaction heats were detected for both these monosaccharides, saturation of binding was not achieved, and as such, values for the enthalpy of binding and binding affinity could not be numerically derived.

Only limited binding of N-acetylglucosamine was observed in soaked crystals (Ziółkowska et al. 2006). A single molecule of N-acetylglucosamine was found in site 6 of the GRFT dimer, whereas other sites were occupied by different ligands present in the mother liquor, such as ethylene glycol (site 5), sulfate (sites 1 and 4), and water (site 2). Site 3 is involved in intermolecular contacts and is not accessible to soaked ligands. Preferential binding of a carbohydrate molecule in site 6 simply indicates easier accessibility, since all six sites are occupied by N-acetylglucosamine in the new cocrystal structure, and the conformation of sugar molecules is similar to that found in the complex with mannose. Additionally, there is a significant difference between the orientation of N-acetylglucosamine in binding site 6 in both the soaked and cocrystal structures (Fig. 2C). The soaked structure was

refined at 2.0 Å resolution, whereas the cocrystal form was refined at 1.56 Å, giving more detailed information and showing clear indications of the presence of both α and β anomers of N-acetylglucosamine in all six binding sites, with the former predominating.

It is worth noting that all sites were fully occupied in the cocrystals although the concentration of N-acetylglucosamine in the crystallization solution was $\sim 15 \text{ mM}$, whereas in the previous experiments the unliganded crystals were soaked in 50 mM solution. Soaking experiments were performed repeatedly with varied concentrations of different ligands and resulted in no carbohydrate binding at lower concentration, and in destruction of the crystals when sugar concentration was raised. Thus the previously reported structure of a complex with N-acetylglucosamine obtained by soaking resulted from an irreproducible experiment in which at least some binding could be seen before crystals were destroyed. Although sulfates occupied some of the carbohydrate-binding sites in the soaked structure, sulfate concentration in the cocrystallization experiments was actually significantly higher. The differences between the structures obtained by soaking versus cocrystallization point out why the latter, if feasible, is by far a preferred method of preparing ligand complexes.

Although the antiviral properties of GRFT are based on its ability to bind efficiently oligosaccharides with branches ending in mannose, the structural data presented here indicate that there is apparently very little difference in the ability of this lectin to bind other monosaccharides, such as glucose and N-acetylglucosamine. The fact that the calorimetric results reported here and those from an earlier study with disaccharides (Ziółkowska et al. 2006) show a thermodynamic preference for mannose and 1 \rightarrow 6 α -mannobiose over glucose and maltose, respectively, indicates




Figure 2. Details of the interactions between glucose and N-acetylglucosamine and GRFT. (A) Glucose in site 1. The location and lengths of hydrogen bonds are representative for all six principal carbohydrate binding sites. (B) N-acetylglucosamine in site 1. Both anomers are present, and the conformation of the side-chain of Asp109 is twofold disordered. (C) Superposition of the molecules of N-acetylglucosamine found in the binding site 6 of GRFT in the cocystal (white) and soaked (cyan) forms of the complexes. Both anomers are present in the cocystal structure, whereas only a single anomer was modeled in the soaked structure.

that further studies are necessary to determine the biophysical source of this preference. Since previously published data (Mori et al. 2005) indicated that mannose was four- to fivefold better than glucose or N-acetylglucosamine at inhibiting the binding of GRFT to HIV gp120, the functional importance of further investigating this dichotomy is clear.

Materials and Methods

GRFT used for structural studies was prepared as recombinant protein in *Nicotiana benthamiana*, expressed and purified as described previously (Ziółkowska et al. 2006). Crystals of the complexes of GRFT with glucose and N-acetylglucosamine

were obtained by the hanging-drop, vapor diffusion method. The crystals of both complexes were grown in 1.8 M MgSO_4 , 0.1 M MES (pH 6.5), with 1:10 molar ratio of GRFT monomers to monosaccharide (protein concentration, ~ 20 mg/mL). The quality of crystals of the complex with glucose was improved by streak seeding. Crystals of both complexes are isomorphous in the space group $P2_12_12_1$ (Table 1), with a dimer in the asymmetric unit.

X-ray diffraction data were collected at the SER-CAT beamline 22-ID, at the Advanced Photon Source (APS), on a MAR 300CCD detector. Prior to data collection, crystals were briefly soaked in a crystallization buffer containing additional 10% ethylene glycol and were rapidly frozen in a nitrogen stream (temperature, 100 K). All data were processed and scaled using the HKL2000 package (Otwinowski and Minor 1997). The structures of both complexes were solved by molecular replacement with PHASER (Storoni et al. 2004) using as a model the coordinates of GRFT dimer from the complex with mannose refined at 1.78 Å resolution (PDB code 2guc) (Ziółkowska et al. 2006). For the complex with glucose, a solution was found with the Z-score of 15.7 after rotation function search and 23.5 after translation function search, and resulted in $R = 0.267$ and $R_{\text{free}} = 0.288$ after a single run of restrained refinement with no rebuilding. The solution for the complex with N-acetylglucosamine was found with the Z-score of 14.3 after rotation function and 27.7 after translation function, and with $R = 0.231$ and $R_{\text{free}} = 0.318$ after the first run of restrained refinement. The structures were further rebuilt with COOT (Emsley and Cowtan 2004) and refined with REFMAC5 (Murshudov et al. 1997). The results of the refinement are summarized in Table 1. The coordinates and structure factors have been deposited in the PDB with accession codes 2nuo and 2nu5 for the experimentally determined complexes of GRFT with glucose and N-acetylglucosamine, respectively.

All calorimetric titration experiments were carried out on a VP-ITC titration calorimeter (Microcal Inc.). In each experiment, 5 μL aliquots of a 1.18 mM solution of a monosaccharide were titrated into a rapidly spinning (300 rpm) solution of a 0.17 mM solution of GRFT (cell volume, 1.3472 mL) held at constant temperature (30°C). A total of 55 injections were made per experiment. All protein solutions were dialyzed in 10 mM PBS (pH 7.4), and the exact concentration of GRFT in each experiment was determined by amino acid analysis on a Hitachi model L-8800 automated amino acid analyzer according to manufacturer's protocols. The binding isotherms were corrected for the heats of dilution, and the resulting area under the




Figure 3. Comparison of the calorimetric heats of interaction of mannose, glucose, and N-acetylglucosamine with GRFT.

Table 2. A comparison of the calorimetric heats of interactions

Monosaccharide	Heats of interaction (cal/mol)
Mannose	-81.11 ± 27.39
Glucose	-50.21 ± 9.80
N-acetylglucosamine	-43.17 ± 22.68

interaction heat peaks were normalized to the amount of monosaccharide and averaged over the course of the experiment.

Acknowledgments

We acknowledge the use of beamline 22-ID of the Southeast Regional Collaborative Access Team (SER-CAT), located at the Advanced Photon Source, Argonne National Laboratory. Use of the APS was supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract No. W-31-109-Eng-38. This project has been funded in whole or in part with federal funds from the National Cancer Institute, National Institutes of Health, under contract N01-CO-12400. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. government. This research was supported in part by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research.

References

- Brünger, A.T. 1992. The free R value: A novel statistical quantity for assessing the accuracy of crystal structures. *Nature* **355**: 472–474.
- Charan, R.D., Munro, M.H., O'Keefe, B.R., Sowder, R.C.I.I., McKee, T.C., Currens, M.J., Pannell, L.K., and Boyd, M.R. 2000. Isolation and characterization of *Myrianthus holstii* lectin, a potent HIV-1 inhibitory protein from the plant *Myrianthus holstii*. *J. Nat. Prod.* **63**: 1170–1174.
- Chothia, C. and Murzin, A.G. 1993. New folds for all- β proteins. *Structure* **1**: 217–222.
- Emsley, P. and Cowtan, K. 2004. Coot: Model-building tools for molecular graphics. *Acta Crystallogr.* **D60**: 2126–2132.
- Mori, T., O'Keefe, B.R., Sowder, R.C., Bringans, S., Gardella, R., Berg, S., Cochran, P., Turpin, J.A., Buckheit Jr., R.W., McMahon, J.B., et al. 2005. Isolation and characterization of griffithsin, a novel HIV-inactivating protein, from the red alga *Griffithsia* sp. *J. Biol. Chem.* **280**: 9345–9353.
- Murshudov, G.N., Vagin, A.A., and Dodson, E.J. 1997. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr.* **D53**: 240–255.
- Otwinowski, Z. and Minor, W. 1997. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol.* **276**: 307–326.
- Storoni, L.C., McCoy, A.J., and Read, R.J. 2004. Likelihood-enhanced fast rotation functions. *Acta Crystallogr.* **D60**: 432–438.
- Ziółkowska, N.E. and Wlodawer, A. 2006. Structural studies of algal lectins with anti-HIV activity. *Acta Biochim. Pol.* **53**: 617–626.
- Ziółkowska, N.E., O'Keefe, B.R., Mori, T., Zhu, C., Giomarelli, B., Vojdani, F., Palmer, K.E., McMahon, J.B., and Wlodawer, A. 2006. Domain-swapped structure of the potent antiviral protein griffithsin and its mode of carbohydrate binding. *Structure* **7**: 1127–1135.
- Ziółkowska, N.E., Shenoy, S.R., O'Keefe, B.R., McMahon, J.B., Palmer, K.E., Dwek, R.A., Wormald, M.R., and Wlodawer, A. 2007. Crystallographic, thermodynamic, and molecular modeling studies of the mode of binding of oligosaccharides to the potent antiviral protein griffithsin. *Proteins* **67**: 661–670.